arXiv:hep-th/9505090v1 17 May 1995

**COMMENT ON: "Integrable Systems in Stringy Gravity"**

This paper [1] is doubtless interesting, unfortunatly almost all the material presented here is already known and has been published elsewhere. Its first part is a standard introduction of generation techniques in General Relativity given even in text books like ref. [2]. The main problem of this generation techniques is to find an explicite expression for the $g \in G$ element of equation (9) in Gal'tsov's paper. This expression is known for some cases like Einstein-Maxwell theory (EM) [3] (for the Einstein-Maxwell plus Dilaton theory $g$ does not modify), n-dimensional gravity in spacetime [4], n-dimensional gravity in potential space [5] [6], etc. (see ref. [7] for a review of this topic). The first step for the solution of this problem in potential space is to find the form of the target space, see equation (15) in Gal'tsov's paper. But this metric is known for the Einstein-Maxwell-Dilaton gravity since 1969, see chapter I in ref. [8], where the metric is written just as in equation (15) of Gal'tsov's paper, but using a parameter $2/(\xi - 3/2)$ instead of $\alpha^2$ of Gal'tsov's. After a Legendre transformation, this metric has been intensively used for finding new exact solutions in five-dimensional gravity (FDG) [9] [10] [11], where $\alpha^2 = 3$ in Gal'tsov's notation, or $\xi = 13/6$ in Neugebauer's notation. Furthermore, this metric has been used for generating exact static solutions in Dilaton gravity for $\alpha$ arbitrary in [12] (using the $SL(2,R)/U(1)$ representation of $g$), where we apply the harmonic map ansatz [13] [14], because we even knew that the target space is non-symmetric for arbitrary $\alpha$, which implies that the Lax pair is useless for the non-static case. In the static case some identifications between EM and FDG can be done, and solutions for one of them can be mapped into the other one, as mentioned at the end of paper [11]. Therefore, the generalization for arbitrary $\alpha$ was trivial for static fields. To see this, it is sufficient to write down the matrix $g \in SL(3,R)$ of equation (4) of ref. [6] for magnetostatic fields, i.e. $\psi = \epsilon = 0$

$$g = \frac{1}{f\kappa^{\frac{2}{3}}} \begin{pmatrix} -2f^2 & 0 & 0 \\ 0 & 2 & -\frac{1}{\sqrt{2}}\chi \\ 0 & -\frac{1}{\sqrt{2}}\chi & -\frac{1}{4}(\chi^2 - f\kappa^2) \end{pmatrix} \qquad (0.1)$$

or for electrostatic fields, i.e. $\chi = \epsilon = 0$

$$g = -\kappa^{\frac{4}{3}} \begin{pmatrix} -2(\psi^2 - \frac{f}{\kappa^2}) & 0 & -\frac{1}{\sqrt{2}}\psi \\ 0 & \frac{2}{f\kappa^2} & 0 \\ -\frac{1}{\sqrt{2}}\psi & 0 & -\frac{1}{4} \end{pmatrix}. \qquad (0.2)$$

After a symmetry transformation, the $SL(2,R)/U(1)$ sectors of matrices (1) and (2) are just the matrix (20) in Gal'tsov's paper, with the following identification given in eq.(7) of ref. [12]

$$\kappa^2 = e^{-2\alpha\phi},$$

and substituting $u = \chi$ for magnetostatic and $u = \psi$ for electrostatic fields. Soliton solutions of the $SL(2,R)/U(1)$ sectors are given in [15] (see also ref. [16]). Nevertheless, we have found that the harmonic map method is more powerful and convenient for this kind of differential equations [12].

Most of the last part of Gal'tsov's paper has been published recently by A. Garcia, O. V. Kechkin and Gal'tsov himself [17] [18]. Again, the target space (23)-(24) of Gal'tsov's paper, and the isometries analysis of the metric in the potential space have been carried out in ref. [18] section III systematically.

To end this comment I must say that although I have used some references of difficult access here, Mr. Gal'tsov cannot argue that he did not know this references. During his stay in Mexico last year (which is surprisingly not mentioned in his paper!!) I personally lend him reference [8], gave him references [6], [10], [16], [11], [12], [7], and he had easy access to the other ones.

I thank Hernando Quevedo for helpful discussions. This work is partially supported by CONACYT-Mexico.

Tonatiuh Matos, Departamento de Física, Centro de Investigación y de Estudios Avanzados del I.P.N., Apdo. Post. 14-740, 07000, México, D.F., México. tmatos@fis.cinvestav.mx.

PACS numbers: 04.50.+h, 11.25.Mj


[1] D. V. Gal'tsov. *Phys. Rev. Lett.* **74**, 2863, (1995).

[2] D. Kramer, H. Stephani, M. MacCallum and E. Held. *Exact Solutions of Einstein Field Equeations.* (1980), DVW, Berlin.

[3] G. Neugebauer and D. Kramer. In *Galaxies, Axisymmetric Systems and Relativity*, Ed. M. A. H. MacCallum.

[4] T. Matos. *Rev. Mex. Fis.* **35**, 208, (1989).

[5] D. Maison. *Phys. Rev. Lett.* **41**, 521, (1978).

[6] T. Matos. *Phys. Lett.* **A131**, 423, (1988).

[7] T. Matos. In: *Proceedings of the International Conference on Aspects of General Relativity and Mathematical Physics*, Ed. N. Breton, R. Capovilla and T. Matos. CINVESTAV, Mexico, (1994).

[8] G. Neugebauer. *Habilitationsschrift* (Universität Jena Press, Jena, Germany) (1969).

[9] T. Matos. *Astron. Nachr.* **307**, 317, (1986).

[10] T. Matos. *Ann. Phys. (Leipzig).* **46**, 462, (1989).

[11] T. Matos. *J. Math. Phys.* **35**, 1302, (1994).

[12] T. Matos, D. Nuñez and H. Quevedo. *Phys. Rev.* **D51**, R310, (1995).

[13] T. Matos and R. Becerril, *Rev. Mex. Fis.* **38**, 69, (1992).

[14] T. Matos. *'Exact Solutions of G-invariant Chiral Equations'* to be published in *Math. Notes* (1995).

[15] T. Matos. *Dissertationschrift* (Universität Jena Press, Jena, Germany) (1987).

[16] D. Kramer, D. Neugebauer and T. Matos. *J. Math. Phys.* **32**, 2727, (1991).

[17] A. Garcia, D. V. Gal'tsov and O. V. Kechkin. *Phys. Rev. Lett.* **74**, 1276, (1995).

[18] D. V. Gal'tsov and O. V. Kechkin. *Phys. Rev.* **D50**, 7394, (1994).